arXiv:hep-ph/9805454v1 26 May 1998

KAIST-TH-96/21

Determination of $|V_{cb}|$ from the polarization of vector meson in the semileptonic decay of B and B_c meson

Myoung-Taek Choi* and Jae Kwan Kim

Department of Physics, Korea Advanced Institute of Science and Technology,
Taejon, 305–701, Korea

(February 1, 2008)

Abstract

Since the degree of polarization of the vector particle in semileptonic decay strongly influences the decay width of the particles, it can be used as a measure of Cabibbo-Kobayashi-Maskawa quark-mixing matrix elements. We show that $|V_{cb}|$ can be determined from the measurement of polarization of vector meson in $\bar{B}, \bar{B}_c \to V l \bar{\nu}$ decay, where V is vector meson.

Typeset using REVTEX

---

*Present address: Physics Dept., Hanyang University, Seoul 133-791, Korea. E-mail: mtchoi@hepth.hanyang.ac.kr

Exclusive semileptonic decays of hadrons containing a bottom quark provide a means to measure the Cabibbo-Kobayashi-Maskawa(CKM) matrix elements [1]. The decay $\bar{B} \to D^* l \bar{\nu}$ is a key process for these studies. Neubert proposed the method of model independent determination of $|V_{cb}|$ from $\bar{B} \to D^* l \bar{\nu}$ decay within the framework of Heavy Quark Effective Theory (HQET) [2]. He observed that the zero recoil point is especially suitable for the extraction of $|V_{cb}|$. His method basically relies on the existence of one universal form factor and the fact that the form factor is unaffected from $1/m_Q$ $(Q = b, c)$ corrections at zero recoil by Luke's theorem [3].

The decays of the $\bar{B}_c$ meson are another source of measurement of CKM element. The $\bar{B}_c \to J/\psi l \bar{\nu} \to (l^+ l^-) l \bar{\nu}$ decay is, in particular, expected to be a quite suitable mode to measure $|V_{cb}|$ due to the experimentally clean signature. Based on the fact that the matrix element of $\bar{B}_c \to J/\psi l \bar{\nu}$ decay is expressed by only one form factor near zero recoil point due to the spin symmetry of heavy mesons, Sanchis *et al.* [4] discussed similar procedure as Neubert to extract $|V_{cb}|$.

In the above method, the value of $|V_{cb}|$ is extracted from the recoil spectrum of the semileptonic decay, hence complete kinematic reconstruction is required. The decay of neutral $\bar{B}$ meson has been used because the flight direction of $\bar{B}^0$ is unaffected by the magnetic field. In ARGUS and CLEO experiments [5–7], where $\bar{B}^0$ is produced almost at rest, the recoil momentum of D^{*+} to the $\bar{B}^0$ meson is easily calculated by reconstructing D^0 and π^+. The problem comes from the small released energy of decay products of D^{*+}; since the momentum of pion of the decay is small, the efficiency for reconstructing the charged pion is low, and the combinatorial background for neutral pions from D^{*0} decays is significant. In ALEPH experiment [8], the $\bar{B}^0$ meson is largely boosted. The momentum of $\bar{\nu}$ is calculated from the missing energy in the hemisphere containing the $D^{*+} l$ candidate. Although the momentum of the pion from the D^{*+} decay has large momentum, the momentum of $\bar{B}^0$ meson should be reconstructed on an event by event basis: the flight direction of the $\bar{B}^0$ meson is measured by reconstructing its production and decay points. There are uncertainties from the missing energy measurement and identification of particles.

In this paper, we show that the polarization of the vector particle can be used as a measure of the CKM element. It is based on the fact that the degree of polarization of the vector particle strongly influences the momentum spectrum of the decay particles as well as the decay rate. One advantage of the polarization measurement is no need of full kinematic reconstruction: the polarization of the vector meson is measured only from their decay products regardless of the charge of the original meson(B or B_c^+).

The differential decay rate for $\bar{B} \to D^* l \bar{\nu}$ is

$$\frac{1}{\sqrt{y^2-1}}\frac{d\Gamma}{dy} = \frac{G_F^2}{48\pi^3} m_{D^*}^3 (m_B - m_{D^*})^2 \eta_{A_1}^2 |V_{cb}|^2 \xi^2(y) \tag{1}$$

$$\times 4y(y+1)\frac{1-2yr+r^2}{(1-r)^2}, \qquad \bar{B} \to D_T^* l \bar{\nu},$$

$$\times (y+1)^2, \qquad \bar{B} \to D_L^* l \bar{\nu}. \tag{2}$$

Here the variable y is

$$y = v \cdot v' = \frac{m_B^2 + m_{D^*}^2 - q^2}{2 m_B m_{D^*}} \tag{3}$$

and $r = m_{D^*}/m_B$, q^2 the momentum transfer to the lepton pair, η_{A_1} the QCD correction factor.

The form factor at non-zero recoil point is not predicted from HQET itself, and can be parameterized by various form. Here we compare three different functional form, linear, exponential, and the pole ansatz [2];

$$\xi(y) = \xi(1)(1 - \bar{\rho}^2(y-1)), \tag{4}$$

$$\xi(y) = \xi(1) e^{-\bar{\rho}^2(y-1)}, \tag{5}$$

$$\xi(y) = \xi(1)(\frac{2}{y+1})^{2\bar{\rho}^2} \tag{6}$$

where $\xi(1) = 1 + \delta(1/m_Q^2)$. Then $|V_{cb}|\xi(v \cdot v')$ is fitted to recoil spectrum with $|V_{cb}|$ and $\bar{\rho}$ as free parameters in the Neubert's method.

The polarization of D^* is measured from the angular distribution of their decay products in the rest frame of D^*. In $\bar{B} \to D^* l \bar{\nu}$ decay where D^* decays into two pseudo-scalar particle D and π, the decay amplitude with the helicity λ of the D^* is given [9]

$$\mathcal{M} \propto \frac{G_F}{\sqrt{2}} V_{cb} \sum_{\lambda} L_\lambda H_\lambda Y^1_\lambda, \tag{7}$$

where $\lambda = 0, \pm 1$ is the helicity of the D^*, and hence also that of the virtual W^- in the limit of massless leptons. L_λ describe the $W^{*-}_\lambda \to l\bar{\nu}$ decay, H_λ are the $\bar{B} \to D^*_\lambda W^{*-}_\lambda$ decay amplitudes and the $J = 1$ spherical harmonics Y^1_λ describe the $D^*_\lambda \to D\pi$ decay.

The angular distribution of π meson in rest frame of D^* is given by

$$\frac{d\Gamma}{d\cos\theta_\pi} \propto 1 + \alpha \cos^2\theta_\pi, \tag{8}$$

where θ_π is the angle between the π direction in the D^* rest frame and the D^* direction in the B rest frame. The polarization parameter α is related to the ratio of longitudinal and transverse decay width;

$$\alpha = 2\,\frac{\Gamma_L}{\Gamma_T} - 1. \tag{9}$$

We note that $|V_{cb}|$ and QCD correction factor are canceled in the Eq. (9). The only unknown parameter is $\bar{\rho}$. Fitting the distribution of D or π in the rest frame of D^* gives the α, and henceforth $\bar{\rho}$. Once the integrated decay rate of Eq.(1) is measured, the value of $|V_{cb}|$ is obtained.

The current measurements of polarization give

$$\alpha = \begin{cases} 0.65 \pm 0.66 \pm 0.25 & [5] \\ 1.48 \pm 0.32 \pm 0.14 & [6] \\ 1.1 \pm 0.4 \pm 0.2 & [7]. \end{cases} \tag{10}$$

Taking the average value of these data, we obtain

$$\alpha = 1.24 \pm 0.25. \tag{11}$$

We show the plot of the $|V_{cb}|$ versus α in Figure 1, using the linear parameterization, Eq (4). We obtain [1]

---

[1] The error in $\bar{\rho}$ is asymmetric. This is due to the functional form of $\bar{\rho}$ with respect to the polarization in Eq. (9). However, the $|V_{cb}|$ is mostly symmetric as seen in Fig. 1.

$$|V_{cb}| = 0.036 \pm 0.005 \tag{12}$$

$$\bar{\rho} = 0.76^{+0.26}_{-0.53}. \tag{13}$$

Different ansatz produce correspondingly slight different values of $\bar{\rho}$ and $|V_{cb}|$. The exponential ansatz leads to $|V_{cb}| = 0.036 \pm 0.006$ and $\bar{\rho} = 0.84^{+0.38}_{-0.60}$, while the pole gives $|V_{cb}| = 0.037 \pm 0.007$ and $\bar{\rho} = 0.89^{+0.40}_{-0.64}$. Here we used [10] $\tau_B = 1.56$ ps, $\mathrm{Br}(\bar{B}^0 \to D^* l \bar{\nu}) = 4.56$ % to obtain $\Gamma_{\bar{B}^0 \to D^* l \bar{\nu}} = 19.2 \times 10^{-15}$ GeV, and $\eta_{A_1}\xi(1) = 0.91$ [11]. The uncertainties of τ_B, $\mathrm{Br}(\bar{B}^0 \to D^* l \bar{\nu})$, and η_{A_1} which are much smaller than that of the polarization are not taken into account in the evaluation of $|V_{cb}|$. We note that the averaged value of $\bar{\rho}$ observed experimentally, based on the fit to the recoil spectrum is $\bar{\rho} = 0.93 \pm 0.06$ [12]. The large error in our method comes from the large experimental uncertainty in the measurement of α.

Now we discuss the decay of $\bar{B}_c$ meson. The $\bar{B}_c \to J/\psi l \bar{\nu}$ decay, followed by $J/\psi \to l^+ l^-$, is the most promising mode to study the semileptonic decay of the $\bar{B}_c$ meson due to its clean signature. Hence the polarization of J/ψ meson might be measured with small experimental uncertainties.

Near zero recoil point the matrix element is expressed only one form factor [4,13];

$$\langle J/\psi, v' | A_\mu | \bar{B}_c, v \rangle = 2\sqrt{m_{B_c} m_{\eta_c}}\ \Delta(v \cdot v')\ \epsilon^*_\mu. \tag{14}$$

where ϵ_μ is the polarization vector of the J/ψ.

In the limit of vanishing lepton mass, one finds;

$$\frac{d\Gamma}{dy} = \frac{G_F^2}{48\pi^3} |V_{cb}|^2 m_{B_c}^2 m_{J/\psi}^3 (y^2 - 1)^{1/2} (y+1)^2 F(r, y) \tag{15}$$

with

$$F(r,y) = 8(1 - 2yr + r^2) \left(\tfrac{\Delta}{y+1}\right)^2, \bar{B}_c \to J/\psi_T\ \mu\ \bar{\nu}_\mu, \tag{16}$$

$$F(r,y) = 4(y - r)^2 \left(\tfrac{\Delta}{y+1}\right)^2, \quad \bar{B}_c \to J/\psi_L\ \mu\ \bar{\nu}_\mu, \tag{17}$$

where the corresponding variable y should be used instead of Eq. (3).

Using operator product expansion, Jenkins *et al.* [13] showed that Δ can be expressed by wave function overlaps of initial and final meson in the zero recoil limit:

$$\Delta(1) = \int d^3\vec{x}\ \Psi^*_{J/\psi}(\vec{x})\ \Psi_{\bar{B}_c}(\vec{x}), \tag{18}$$

which is the wave function overlap of initial and final hadron. Because the flavour symmetry is broken, $\Delta(v \cdot v')$ is not normalized to 1 at zero recoil point. The wave function overlap might be calculated from the lattice QCD calculation or quark potential models in a reliable way because of non-relativistic nature of the $\bar{B}_c$ meson.

Deviations from the spin symmetry imply the appearance of new form factors in the hadronic matrix element which do not contribute at zero recoil. Spin symmetry breaking effects can occur when the c-quark's recoil momentum is larger than m_c. However we expect that the Eq. (14) is applicable to other kinematic point since the recoil momentum of J/ψ is small ($y_{\text{max}} - 1 \simeq 0.26$) due to its heavy mass.

In $\bar{B}_c \to J/\psi l \bar{\nu}$ decay where the J/ψ meson decays into two lepton pair, the decay amplitude with the helicity λ of the J/ψ meson is obtained in a similar way to Eq. (7), corresponding Y^1_λ be replaced by Wigner d-function

$$\mathcal{M} \propto \frac{G_F}{\sqrt{2}} V_{cb} \sum_\lambda L_\lambda H_\lambda d^1_{\lambda_{J/\psi}, \lambda_{l^-} - \lambda_{l^+}}, \tag{19}$$

where $\lambda_{l^-} - \lambda_{l^+} = \pm 1$.

The angle distribution of l^- in rest frame of J/ψ is given

$$\frac{d\Gamma}{d\cos\theta_{l^-}} \propto 1 + \alpha' \cos^2\theta_{l^-}, \tag{20}$$

where we find

$$\alpha' = \frac{\Gamma_T - 2\Gamma_L}{\Gamma_T + 2\Gamma_L}. \tag{21}$$

The parameter α' is again related to the ratio of longitudinal and transverse decay widths. With the observation that the $\Delta(1)$ is canceled from the expression, one obtains $\bar{\rho}$ from the polarization of the J/ψ meson. Since unknown two parameters are obtained [2], one can calculate $|V_{cb}|$ once the integrated decay width is measured, as in $B \rightarrow D^* l \bar{\nu}$ case.

In summary, we discussed an alternate way to extract $|V_{cb}|$ from the measurement of polarization of vector meson in the exclusive semileptonic decays. The advantage of the method is that the full kinematic reconstruction is not needed in this case.

Full use of kinematic information such as angular correlations of the decay products is expected to allow a determination of the invariant hadronic form factors and the $|V_{cb}|$ quark mixing matrix element almost model-independently [9]. The semileptonic decays of B_c meson is quite suitable for those study due to the clean signature.

Acknowledgements

This work was supported in part by the Korea Science and Engineering Foundation.

---

[2] We note that one should be careful to parameterize the form factor $\Delta(y)$ in the linear form like Eq.(4) because $\bar{\rho}^2$ is predicted to be large, even $2 \sim 4$ in ISGW model [14]

REFERENCES

[1] M. Kobayashi and T. Maskawa, Prog. Theo. Phys. **49** (1973) 652; N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531.

[2] M. Neubert, Phys. Lett. B **264** (1991) 455.

[3] M. Luke, Phys. Lett., **B252** (1990) 447.

[4] M. A. Sanchis, Nucl. Phys. **B440** (1995) 251; Phys. Lett. **B312** (1993) 333.

[5] ARGUS Collaboration, H. Albrecht *et al.,* Phys. Lett. **219** (1989) 121; Z. Phys. **C57** (1993) 523; Phys. Lett. **B324** (1994) 249; **275** (1992) 175.

[6] CLEO Collaboration, D. Bortoletto *et al.*, Phys. Rev. Lett. **63** (1989) 1667; R. Fulton *et al.*, Phys. Rev. **D43** (1991) 651; B. Barish *et al.*, Phys. Rev. **D51** (1995) 1014.

[7] CLEO Collaboration, A. Ryd *et al.*, in *The Albuquerque Meeting*, Proceedings of the 8th Meeting of the Division of Particles and Fields of the American Physical Society, Albuquerque, 1994, edited by Sally Seidel (World Scientific, Singapore, 1995), p. 847; S. Sangheera *et al.*, Phys. Rev. **D47** (1993) 791.

[8] ALEPH Collaboration, D. Buskulic *et al.*, Phys. Lett. **B359** (1995) 236.

[9] K. Hagiwara, A. Martin and M. Wade, Nucl. Phys. **B327** (1989) 596; K. Hagiwara, A. Martin and M. Wade, Phys. Lett. **B228** (1989) 144.

[10] Particle Data Group, Phys. Rev. **D54** (1996) 1.

[11] M. Neubert, Nucl. Phys. **B371** (1992) 149.

[12] T.E. Browder (CLEO Collaboration), in: Proc. of the XXVII Int. Conf. on High Energy Physics, Glasgow, Scotland, July 1994, eds. P.J. Bussey and I.G. Knowles (IOP Publ. Ltd., Bristol, 1995), Vol. 2, p. 1117; I.J. Scott (ALEPH Collaboration), *ibid.* Vol. 2, p. 1121; H. Albrecht et al. (ARGUS Collaboration), Z. Phys. C **57** (1993) 533.

[13] E. Jenkins, M. Luke, A.V. Manohar and M.J. Savage, Nucl. Phys. **B390** (1993) 463.

[14] N. Isgur, D. Scora, B. Grinstein and M.B. Wise, Phys. Rev. **D39** (1989) 799.

FIGURES

FIG. 1. $|V_{cb}|$ versus α in $\bar{B} \to D^* l \bar{\nu}$ (using the linear parametrization of the form factor). $\tau_B = 1.56 \pm 0.06$ ps, $\mathrm{Br}(\bar{B} \to D^* l \bar{\nu}) = 4.56 \pm 0.27$ % is used



|V_cb|
0.044
0.042
0.04
0.038
0.036
0.034
0.032
0.03
0.9
1
1.1
1.2
1.3
1.4
1.5
α